

## GENETIC AND PHYLOGENETIC CHARACTERIZATION OF RABIES VIRUS ISOLATES FROM WILDLIFE AND LIVESTOCK IN PARAIBA, BRAZIL

Y. SHOJI<sup>1</sup>, Y. KOBAYASHI<sup>1</sup>, G. SATO<sup>1</sup>, A.A.B. GOMES<sup>2</sup>, T. ITOU<sup>1\*</sup>, F.H. ITO<sup>3</sup>, T. SAKAI<sup>1</sup>

<sup>1</sup>Department of Preventive Veterinary Medicine and Animal Health, School of Veterinary Medicine, Nihon University, 1866 Kameino, Fujisawa, Kanagawa 252-8510, Japan; <sup>2</sup>Department of Veterinary Medicine, Center of Health and Rural Technology, Federal University of Campina Grande, Patos, Paraiba, Brazil; <sup>3</sup>Department of Preventive Veterinary Medicine and Animal Health, Faculty of Veterinary Medicine and Zootechny, University of Sao Paulo, Sao Paulo, Brazil

Received April 25, 2005; accepted December 20, 2005

**Summary.** – Thirty-four rabies virus (RV) isolates from foxes (8), insectivore bats (9), cattle (14), sheep (1), a goat (1) and a donkey (1) from Paraiba state, northeastern Brazil, were genetically characterized. Sequences of 890 nts of nucleoprotein (N) genes of these isolates were analyzed and compared with those of other Brazilian isolates characterized earlier. Phylogenetic analysis revealed three genetical lineages of RV co-existing in this region. Each lineage was found to be associated with particular host species and to circulate independently of each other. The first lineage was found in foxes (*Dusicyon* sp.) and could be discriminated from domestic carnivore isolates from Sao Paulo, Goias and Minas Gerais in the southern and central Brazil. The second lineage was associated with insectivorous bats (*Molossus* spp.) and differed from vampire bat-associated RV isolates. The third lineage was found in livestock and clustered with vampire bat-associated RV isolates from Sao Paulo, Tocantins, Goias and Matto Grosso. These results indicate that RV of these genetic lineages are co-circulating in the Paraiba state and that livestock in this region are infected with vampire bat-associated RV, suggesting that the vampire bat is the main reservoir of livestock rabies in this region.

**Key words:** Brazil; domestic animals; epidemiology; phylogenetic analysis; Rabies virus; wild animals

### Introduction

Rabies is an enzootic disease caused by RV, a neurotropic virus of the genus *Lyssavirus*. Lyssaviruses (members of the *Lyssavirus* genus) are divided in 7 genotypes (GTs), among them GT1 (RV), GT3 (Mokola virus, MOKV) and GT7 (Australian bat lyssavirus, ABLV) (Fauquet *et al.*, 2005).

RV infects almost all kinds of mammalian species. RV infects the central nervous system, causing acute encephalitis, which is almost incurable after the clinical symptoms appear. However, there is a possibility of

prevention of this disease by effective vaccine even when administered after exposure to the virus.

In Brazil, like most other South American countries, rabies occurs in two different epidemiological forms, sylvatic and urban. The main reservoir of urban rabies is the dogs, while that of sylvatic rabies is the vampire bat (Ito *et al.*, 2001a). In large cities of southern Brazil like Rio de Janeiro and Sao Paulo, dog rabies has been controlled by effective dog vaccination campaigns (Romijn *et al.*, 2003; Heinemann *et al.*, 2002). However, in rural areas, the dog remains the main reservoir of human rabies, and the rabies cases of humans and livestock contracted from wild animals, such as vampire bats (*Desmodus rotundus*) (Schneider *et al.*, 1996; Romijn *et al.*, 2003), marmosets (Favoretto *et al.*, 2001), non-human primates (Ramos *et al.*, 2002) and foxes have been reported as increasing (Almeida *et al.*, 2001), especially in the northern Brazil. In the Ceara state, the northeastern

\*E-mail: takuya@brs.nihon-u.ac.jp; fax: +81466-843380.

**Abbreviations:** ABLV = Australian bat lyssavirus; GT = genotype; MOKV = Mokola virus; N = nucleoprotein; RV = Rabies virus

Brazil, 13 human rabies cases originated from wild animals, such as bats, crab-eating raccoons (*Procyon cancrivorus*) and white-tufted-ear marmosets (*Calithrix jacchus jacchus*) during the past 7 years (1991–1998) (Favoretto *et al.*, 2001). The latter study revealed that these viruses represent a unique and independent rabies endemic cycle. In some other states surrounding Paraiba state, outbreaks of rabies in humans transmitted by vampire bats (Bahia, Amazonas and Para states) (Goncalves *et al.*, 2002; Schneider *et al.*, 2001; Pan American Health Organization, 2004) also occurred.

Although rabies cases in humans and livestock in the Paraiba state have occurred, there is so far no report on molecular analysis of the respective RV isolates. The aim of this study was to characterize genetically RV circulating in this area and to obtain epidemiological information for prediction and prevention of rabies outbreaks in both humans and livestock in this region.

### Materials and Methods

**Viruses.** Thirty-four RV isolates obtained from brain specimens from foxes (8), cattle (14), insectivorous bats (9), sheep (1), a goat (1) and a donkey (1), collected in the Paraiba state, the northeastern Brazil, were examined.

**RT-PCR.** Total RNA was extracted from brain specimens using a QIAamp Viral RNA Mini Kit (QIAGEN) according to the manufacturer's instructions. To detect RV RNA in general, a RT-PCR specific for RV N gene (a sequence of 964 nts) was performed using the primer pair P1/P2. To detect carnivore-associated RV RNA in particular, the primer pair RHN1/RHNS3 yielding a product of 1,512 nts was used (Ito *et al.*, 1999, 2003).

**Sequencing.** The RT-PCR products of 964 nts were purified using a QIAquick PCR purification Kit (QIAGEN) and subjected to direct sequencing using the primers P1, P2, BRABN-S1, BRABN-S3 and BRABN-C3 (Shoji *et al.*, 2004), a Big Dye Terminator Cycle Sequencing Kit version 3.1 (Applied Biosystems) and an automated sequencer (Applied Biosystems).

**Phylogenetic analysis** of RV isolates was based on the alignment of 890 nts-long sequences of N gene (nt 89–978) (Tordo *et al.*, 1986) using the CLUSTAL X program (Thompson *et al.*, 1994). A neighbor-joining tree was constructed using a method of Saitou and Nei (1987) and the bootstrap probabilities of each node were calculated using 1,000 replicates. A graphic output was obtained using the TREEVIEW program (Page *et al.*, 1996).

### Results

#### *Detection of RV RNA by RT-PCR*

All the 34 RV isolates were positive by the RT-PCR with the primer pair P1/P2, while only 8 of them, originating exclusively from fox brains, were positive by the RT-PCR

with the primer pair RHN1/RHNS3 (Fig. 1); the latter isolates thus represented carnivore-associated RV.

#### *Sequencing and phylogenetic analysis of RV isolates*

The sequences of 890 nt-long RT-PCR products (nt 89–978) corresponding to RV N gene were determined for all the 34 RV isolates and were deposited in the Gen Bank database (Acc. Nos. AB206407–AB206439 and AB207884).

The nucleotide sequence identities of all the isolates with RV (GT1), MOKV (GT3) and ABLV (GT7) were 81.7–99.4%, 72.2–72.7% and 75.9–78.9%, respectively, showing that these isolates belong to GT 1 of lyssaviruses. The nucleotide and deduced amino acid sequences exhibited different levels of identity with each other, 82.5–100% and 92.3–100%, respectively. Among all the fox isolates, the nucleotide sequences showed a 94.6–100% identity. The latter also showed the highest identity (93.2–94.6%) with those of domestic carnivore isolates from Sao Paulo, Minas Gerais and Goias in the southern and central Brazil. However, these levels of identity were lower than those (98.9–99.1%) within domestic carnivore isolates except for fox isolates.

The nucleotide sequence identity within the insectivorous bat isolates from Paraiba was 97.8–99.7%, while the nucleotide and deduced amino acid sequences identities of these isolates with other isolates described in this study were relatively low, 82.1–89.5% and 92.9–97.3%, respectively. The nucleotide sequence identity within the livestock isolates showed 98.2–100%. The nucleotide and amino acid identities of the livestock isolates from Paraiba with the isolates from vampire bats and livestock from other regions in Brazil (Goias, Mato Grosso, Sao Paulo and Tocantins) were the highest, 96.1–98.2% and 97.6–100%, respectively.

Multiple alignment of deduced amino acid sequences (298 aa) of representative RV isolates revealed 19 amino acid substitutions. Several of them appeared to be lineage-specific. Five substitutions were found in carnivore or bat isolates. Another seven substitutions resulted in an amino acid with changed chemical properties. Seven substitutions were found in insectivorous bat isolates (data not shown).

The phylogenetic analysis based on the sequences of 890 nts of N gene revealed that 55 isolates analyzed in this study clustered into four major genetic lineages: (A) the lineage of domestic carnivore isolates from southern and central Brazil; (B) the lineage of fox isolates from Paraiba; (C) the lineage of insectivorous bat isolates from Paraiba and Sao Paulo; (D) the lineage of livestock isolates from Paraiba and vampire bat and livestock isolates from southern and central Brazil (Fig. 2a). Insectivorous bat isolates from Paraiba together with BRNL-1 constituted a lineage that differed from other insectivorous bat isolates from Sao Paulo (BREF-4 and BRMA-1). Fox isolates from Paraiba formed two discrete sub-lineages (B-1 and B-2) and each sub-lineage




Fig. 1

#### Discrimination of carnivore-associated and non-carnivore-associated RV isolates by RT-PCR

Positive controls: BRdg 2 (Brazilian carnivore-associated RV, lanes 1) and BRvmbt 33 (Brazilian non-carnivore RV, lanes 3). Brazilian RV isolates BR-Pfx1 (lanes 2), BR-Pbt 1 (lanes 4) and BR-Pbv 1 (lanes 5). Negative control: RT-PCR without RNA template (lanes 6). P and R indicate P1/P2 and RHN1/RHNS3 primer pairs, respectively.

was distinguishable from domestic carnivore isolates from southern and central Brazil. Although the lineages A and B were categorized in carnivore rabies, they did not contain carnivore isolates from other countries (Fig. 2b). However, the lineages C and D were assigned to Chiropteran rabies together with other bat isolates. The insectivorous bat lineage C has closer relationship with North American and other Brazilian insectivorous bat isolates than the vampire bat lineage.

#### Discussion

In this study, viral RNA was detected by RT-PCR in 34 brain specimens from wild animals (foxes and insectivorous bats), livestock (cattle, sheep, a goat and a donkey) from Paraiba, northeastern Brazil. N gene sequences of these isolates and their phylogenetic analysis revealed that three genetic lineages of RV associated with specific host species occur in this area, indicating the presence of multiple endemic cycles of rabies. The fox RV lineage from Paraiba was found to belong to the group of Brazilian carnivore RV isolates but it apparently differed from carnivore RV lineage from southern and central Brazil. The insectivorous bat (*Molossus* sp.)-RV lineage of Paraiba clustered with an isolate from another species of insectivorous bat (*Nictinomops laticaudatus*; BRNL-1) and could be differentiated from other Brazilian RV isolates. Furthermore, the livestock RV lineage from Paraiba was found to belong to Brazilian vampire bat-associated RV lineages, suggesting that the main reservoir of livestock rabies in Paraiba is vampire bats though we could not isolate RV from vampire bats captured in Paraiba. The high bootstrap values of the node of each

lineage in the phylogenetic tree indicate that these three RV lineages have been circulating independently of each other in this region.

Considering the population of the fox *Dusysion* sp. in Paraiba and the fact that some residents keep foxes as pets, this species has a potential for transmission of RV to humans.

Vampire bats are responsible for most cases of livestock rabies and some outbreaks of human rabies. Fortunately, the human rabies cases transmitted by insectivorous bats have not been reported recently in Brazil, but the growing incidence of humans bitten by bats including insectivorous bats in Brazil (Mayen *et al.*, 2003) and the growing incidence of human rabies transmitted by insectivorous bats in North America (Messenger *et al.*, 2002; Pape *et al.*, 1999) suggest the importance of insectivorous bats in the epidemiology of rabies in Brazil.

In accord with problem of grouping of RV isolates worldwide, Brazilian RV isolates have been divided in two major phylogenetical groups, the carnivore RV group and the Chiropteran RV group (Tordo *et al.*, 1986). The Brazilian RV isolates from foxes, a dog and a cat belong to the carnivore RV group, while the bat and livestock isolates belong to the Chiropteran RV group. In our phylogenetic analysis, the fox, dog and cat, and insect bat RV lineages formed separate clusters without any isolates from other countries, suggesting that each lineage is circulating in Brazil only. On the other hand, the vampire bat RV lineage clustered with vampire bat isolates from other South American countries.

Brazil is so much diverse in geomorphology and in mammalian fauna including bats (Findley, 1993; Walker, 2001) that it is difficult to completely comprehend or elucidate the




Fig. 2

## Phylogenetic analyses of RV isolates based on 890 nt-long sequences of N gene

(a) Phylogenetic tree of Brazilian RV isolates. (b) Phylogenetic tree of representative RV isolates from Brazil and other countries. MOKV (Mokola) and ABLV (ABL) (Acc. Nos. Y09762 and AF418014, respectively) were used as outgroups. Isolates from Paraiba are in bold. A, B (B-1 and B-2), C and D represent individual lineages (sublineages). Percentage bootstrap values out of 1,000 replicates are indicated at each respective node.

ecology of RV in Brazil. The findings of this study in agreement with those of other studies (Favoretto *et al.*, 2001, 2002; Ito *et al.*, 2001b; Romijn *et al.*, 2003) indicate that the RV lineages in Paraiba exhibit a geographic area-dependent divergence in addition to host species-specific divergence, and that endemic cycles of the carnivore-associated RV lineage and the bat-associated RV lineage co-exist in one area. Further investigation in each area and exchanging of epidemiological information is necessary for better understanding and more effective prevention of rabies outbreaks.

**Acknowledgement.** This work was supported in part by the Academic Frontier Project for Private Universities from the Ministry of Education, Culture, Sports, Science and Technology (MEXT) of Japan.

### References

- Almeida MF, Massad E, Aguiar EAC, Martorelli LFA, Joppert AMS (2001): Neutralizing antirabies antibodies in urban terrestrial wildlife. *J. Wildl. Dis.* **37**, 394–398.
- Arai YT, Kuzmin IV, Kameoka Y, Botvinkin AD (2003): New lyssavirus genotype from the lesser mouse-eared bat (*Myotis blythi*), Kyrgyzstan. *Emerg. Infect. Dis.* **9**, 333–336.
- Fauquet CM, Mayo MA, Maniloff J, Dessellberger U, Ball LA (Eds) (2005): *Virus Taxonomy. Eighth Report of the International Committee on Taxonomy of Viruses.* Elsevier-Academic Press, Amsterdam.
- Favoretto SR, Carrieri ML, Cunha EMS, Aguiar EAC, Silva LHQ, Sodew MM, Souza M, Kotaito I (2002): Antigenic typing of Brazilian rabies virus samples isolated from animals and humans, 1989–2000. *Rev. Inst. Med. Trop. Sao Paulo* **44**, 91–95.
- Favoretto SR, de Mattos CC, Morais NB, Alves Araujo FA, de Mattos CA (2001): Rabies in marmosets (*Callithrix jacchus*), Ceara, Brazil. *Emerg. Infect. Dis.* **7**, 1062–1065.
- Findley JS (1993): *Bats: a community perspective.* Cambridge University Press, Cambridge.
- Goncalves MAS, Sa-Neto RJ, Brazil TK (2002): Out break of aggressions and transmission of rabies in human beings by vampire bats in northeastern Brazil. *Rev. Soc. Bras. Med. Trop.* **35**, 462–464.
- Heinemann MB, Fernandes-Matioli FMC, Cortez A, Soares RM, Sakamoto SM, Bernardi F, Ito FH, Madeira AMBN, Richtzenhain LJ (2002): Genealogical analyses of rabies virus strains from Brazil based on N gene alleles. *Epidemiol. Infect.* **128**, 503–511.
- Ito M, Arai YT, Itou T, Sakai T, Ito FH, Takasaki T, Kurane I (2001a): Genetic characterization and geographic distribution of rabies virus isolates in Brazil: identification of two reservoirs, dogs and vampire bats. *Virology* **284**, 214–222.
- Ito M, Itou T, Sakai T, Santos FCM, Arai YT, Takasaki T, Kurane I, Ito FH (2001b): Detection of rabies virus RNA isolated from several species of animals in Brazil by RT-PCR. *J. Vet. Med. Sci.* **63**, 1309–1313.
- Ito M, Itou T, Shoji Y, Sakai T, Ito FH, Arai YT, Takasaki T, Kurane I (2003): Discrimination between dog-related and vampire bat-related rabies viruses in Brazil by strain-specific reverse transcriptase-polymerase chain reaction and restriction fragment length polymorphism analysis. *J. Clin. Virol.* **26**, 317–330.
- Ito N, Sugiyama M, Oraveerakul K, Piyaviriyakul P, Lumlerdacha B, Arai YT, Tamura Y, Mori Y, Minamoto N (1999): Molecular epidemiology of rabies in Thailand. *Microbiol. Immunol.* **43**, 551–559.
- Messenger SL, Smith JS, Rupprecht C (2002): Emerging epidemiology of bat-associated cryptic cases of rabies in humans in the United States. *Emerg. Infect. Dis.* **35**, 738–747.
- Page RD (1996): TreeView: an application to display phylogenetic trees on personal computers. *Comput. Appl. Biosci.* **12**, 357–358.
- Pan American Health Organization (2004): Human rabies transmitted by bats in Para state, Brazil. Emerging and Reemerging Infectious diseases, Region of the Americas. *EID weekly updates* **2**(13)-2 April.
- Pape WJ, Fitzsimmons TD, Hoffman RE (1999): Risk for rabies transmission from encounters with bats, Colorado, 1977–1996. *Emerg. Infect. Dis.* **5**, 433–437.
- Ramos PM, Ramos PS (2002): Human accidents with monkeys relation to prophylactic treatment for rabies, in the municipal district of Sao Paulo, Brazil. *Rev. Soc. Bras. Med. Trop.* **35**, 575–577.
- Romijn PC, Van Der Heide R, Cattaneo CA, Silva RD, Van Der Poel WH (2003): Study of lyssaviruses of bat origin as a source of rabies for other animal species in the state of Rio de Janeiro, Brazil. *Am. J. Trop. Med. Hyg.* **69**, 81–86.
- Saitou N, Nei M (1987): The neighbor-joining method: A new method for reconstructing phylogenetic trees. *Mol. Biol. Evol.* **4**, 406–425.
- Schneider MC, Aron J, Santos-Burgoa C, Uieda W, Ruiz-Velazco S (2001): Common vampire bat attacks on humans in Village of the Amazon region of Brazil. *Cad. Saude Publica* **17**, 1531–1536.
- Schneider MC, Santos-Burgoa C, Aron J, Munoz B, Ruiz-Velazco S, Uieda W (1996): Potential force of infection of human rabies transmitted by vampire bats in the Amazonian region of Brazil. *Am. J. Trop. Med. Hyg.* **55**, 680–684.
- Shoji Y, Kobayashi Y, Sato G, Itou T, Miura Y, Mikami T, Cunha E, Samara S, Carvalho A, Nocitti D, Ito FH, Kurane I, Sakai T (2004): Genetic characterization of rabies viruses isolated from frugivorous bat (*Artibeus* spp.) in Brazil. *J. Vet. Med. Sci.* **66**, 1271–1273.
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997): The CLUSTAL-X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* **25**, 4876–4882.
- Tordo N, Poch O, Ermine A, Keith G, Rougeon F (1986): Walking along the rabies genome: is the large G-L intergenic region a remnant gene? *Proc. Natl. Acad. Sci. USA* **83**, 3914–3918.